

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Topology and its Applications 153 (2006) 2026–2034

---

**Topology  
and its  
Applications**


---

[www.elsevier.com/locate/topol](http://www.elsevier.com/locate/topol)

# Embeddings of homology equivalent manifolds with boundary

D. Gonçalves<sup>a,1</sup>, A. Skopenkov<sup>b,\*</sup>

<sup>a</sup> *Departamento de Matemática, IME, University of São Paulo, Caixa Postal 66281,  
Agência Cidade de São Paulo 05311-970, São Paulo, SP, Brazil*

<sup>b</sup> *Department of Differential Geometry, Faculty of Mechanics and Mathematics, Moscow State University,  
Moscow, Russia 119992*

Received 23 January 2005; received in revised form 3 July 2005; accepted 22 July 2005

---

## Abstract

We prove a theorem on equivariant maps implying the following two corollaries:

(1) Let  $N$  and  $M$  be compact orientable  $n$ -manifolds with boundaries such that  $M \subset N$ , the inclusion  $M \rightarrow N$  induces an isomorphism in integral cohomology, both  $M$  and  $N$  have  $(n - d - 1)$ -dimensional spines and  $m \geq \max\{n + 3, \frac{3n+2-d}{2}\}$ . Then the restriction-induced map  $\text{Emb}^m(N) \rightarrow \text{Emb}^m(M)$  is bijective. Here  $\text{Emb}^m(X)$  is the set of embeddings  $X \rightarrow \mathbb{R}^m$  up to isotopy (in the PL or smooth category).

(2) For a 3-manifold  $N$  with boundary whose integral homology groups are trivial and such that  $N \not\cong D^3$  (or for its special 2-spine  $N$ ) there exists an equivariant map  $\tilde{N} \rightarrow S^2$ , although  $N$  does not embed into  $\mathbb{R}^3$ .

The second corollary completes the answer to the following question: for which pairs  $(m, n)$  for each  $n$ -polyhedron  $N$  the existence of an equivariant map  $\tilde{N} \rightarrow S^{m-1}$  implies embeddability of  $N$  into  $\mathbb{R}^m$ ? An answer was known for each pair  $(m, n)$  except  $(3, 3)$  and  $(3, 2)$ .

© 2005 Published by Elsevier B.V.

---

\* Corresponding author.

E-mail addresses: [dlgoncal@ime.usp.br](mailto:dlgoncal@ime.usp.br) (D. Gonçalves), [skopenko@mccme.ru](mailto:skopenko@mccme.ru) (A. Skopenkov).

<sup>1</sup> Gonçalves supported in part by FAPESP, Projeto Temático Topologia Algébrica, Geométrica e Diferencial-2000/05385-8. Skopenkov supported in part by the Russian Foundation for Basic Research, Grants No. 02-01-00014 and No. 01-01-00583, President of Russian Federation Grant for Support of Scientific Schools NSH-1988.2003.1, Moscow State University Stipendium for Young Scientists, Russian Academy of Sciences Program 'Modern problems of theoretical mathematics' and INTAS Grant No. YSF-2002-393.

MSC: primary 57Q35, 57R40; secondary 55S15, 57Q30, 57Q60, 57Q65

Keywords: Embedding; Deleted product; Homology equivalence; Local coefficients; Homology ball

This paper is on the classical problem of classification of embeddings into Euclidean spaces, and as a main tool we use the Haefliger–Wu invariant. We begin with the formulation of our main homotopy result. Let  $\tilde{N} = \{(x, y) \in N \times N \mid x \neq y\}$  be the *deleted product* of  $N$ . Let  $\mathbb{Z}_2$  act on  $\tilde{N}$  and on  $S^{m-1}$  by exchanging factors and antipodes, respectively. Denote by  $\pi_{\text{eq}}^{m-1}(\tilde{N})$  be the set of equivariant maps  $\tilde{N} \rightarrow S^{m-1}$  up to equivariant homotopy. The set  $\pi_{\text{eq}}^{m-1}(\tilde{N})$  can be effectively calculated [4, beginning of §2], [6], [7, 1.7.1], [20], [1, 7.1], [19, §6]. Note that  $\pi_{\text{eq}}^{m-1}(\tilde{N}) = \emptyset$  for  $m < n$  because  $\tilde{N} \supset \tilde{D}^n \simeq_{\text{eq}} S^{n-1}$ .

**Theorem.** *Let  $N$  and  $M$  be compact orientable connected  $n$ -manifolds with non-empty boundaries such that  $M \subset N$  and the inclusion  $M \rightarrow N$  induces an isomorphism in cohomology with  $\mathbb{Z}$ -coefficients. Then the restriction-induced map  $\pi_{\text{eq}}^{m-1}(\tilde{N}) \rightarrow \pi_{\text{eq}}^{m-1}(\tilde{M})$  is bijective.*

*More precisely, if the inclusion-induced homomorphism  $H^i(N) \rightarrow H^i(M)$  is an isomorphism only for  $i \geq l > 0$ , then the above restriction-induced map is bijective for  $m \geq n + l$  and surjective for  $m = n + l - 1$ .*

This algebraic result is interesting because of the following geometric corollaries. Denote  $\text{CAT} = \text{DIFF}$  or  $\text{PL}$ . For a CAT manifold  $N$  let  $\text{Emb}_{\text{CAT}}^m(N)$  be the set of CAT embeddings  $N \rightarrow \mathbb{R}^m$  up to CAT isotopy. A folklore general conjecture, supported by many known results (for a survey see, e.g., [12, 20]) is that  $\text{Emb}_{\text{CAT}}^m(N)$  is not changed under homology equivalence of  $N$  (i.e., under a map  $f : M \rightarrow N$  between manifolds inducing an isomorphism in (co)homology), in the PL case for  $m \geq n + 3$  and in the DIFF case for  $m \geq \frac{3n}{2} + 2$ .

**Corollary.** *Let  $N$  and  $M$  be compact orientable  $n$ -manifolds with non-empty boundaries such that  $M \subset N$ , the inclusion  $M \rightarrow N$  induces an isomorphism in cohomology with  $\mathbb{Z}$ -coefficients, both  $M$  and  $N$  has  $(n - d - 1)$ -dimensional spine and  $m \geq \max\{n + 3, \frac{3n+2-d}{2}\}$ . Then the restriction-induced map  $\text{Emb}_{\text{CAT}}^m(N) \rightarrow \text{Emb}_{\text{CAT}}^m(M)$  is bijective.*

*More precisely, if the inclusion-induced homomorphism  $H^i(N) \rightarrow H^i(M)$  is an isomorphism only for  $i \geq l > 0$ , then the above restriction-induced map is bijective for  $m \geq n + l$  and surjective for  $m = n + l - 1$ .*

In the DIFF category the restriction  $m \geq \frac{3n+2-d}{2}$  can be relaxed to  $m \geq \frac{3n+1-d}{2}$ .

By the Corollary, any homology ball unknots in codimension at least 3 (cf. [14]).

Recall that a subpolyhedron  $K$  of a manifold  $N$  is called a *spine* of  $N$  if  $N$  is a regular neighborhood of  $K$  in  $N$  (or, equivalently, if  $N$  collapses to  $K$ ) [13]. We remark that for a compact connected  $n$ -manifold  $N$  with boundary, the property of having an  $(n - d - 1)$ -dimensional spine is close to  $d$ -connectedness of  $(N, \partial N)$ . Indeed, for a compact connected  $n$ -manifold  $N$  with boundary and an  $(n - d - 1)$ -dimensional spine, the pair  $(N, \partial N)$  is homologically  $d$ -connected. On the other hand, every compact connected

$n$ -manifold  $N$  with boundary for which  $(N, \partial N)$  is  $d$ -connected,  $\pi_1(\partial N) = 0$ ,  $d + 3 \leq n$  and  $(n, d) \notin \{(5, 2), (4, 1)\}$ , has an  $(n - d - 1)$ -dimensional spine [22, Theorem 5.5], [8, Lemma 5.1 and Remark 5.2]. Recall that a closed manifold  $N$  (or a pair  $(N, \partial N)$ ) is called *homologically  $d$ -connected*, if  $N$  is connected and  $H_i(N) = 0$  for each  $i = 1, \dots, d$  (or  $H_i(N, \partial N) = 0$  for each  $i = 0, \dots, k$ ).

The Corollary follows from the Theorem and the bijectivity of  $\alpha$ -invariant [7, 12, §4], [19, Theorems 1.1 $\alpha\partial$  and 1.3 $\alpha\partial$ ], which is defined as follows. For an embedding  $f: N \rightarrow \mathbb{R}^m$  define a map

$$\tilde{f}: \tilde{N} \rightarrow S^{m-1} \quad \text{by} \quad \tilde{f}(x, y) = \frac{fx - fy}{|fx - fy|}.$$

The equivariant homotopy class  $\alpha(f)$  of the above-defined  $\tilde{f}$  in  $\pi_{\text{eq}}^{m-1}(\tilde{N})$  is clearly an isotopy invariant. Thus is defined the *Haefliger–Wu (deleted product) invariant*

$$\alpha = \alpha_{\text{CAT}}^m(N) : \text{Emb}_{\text{CAT}}^m(N) \rightarrow \pi_{\text{eq}}^{m-1}(\tilde{N}).$$

We remark that the assumption that  $(N, M)$  is a codimension 0 pair is essential in the Theorem and the Corollary. Indeed, take  $N = D^p \times S^q$  and  $M = S^q$ . For  $m \geq 3q/2 + 2$  we have  $\#\text{Emb}^m(S^q) = 1$  while  $\text{Emb}^m(D^p \times S^q) = \pi_q(V_{m-q,p})$  can contain more than one element (specific examples are particularly easy to find for  $p = 1$ , when  $V_{m-q,p} \simeq S^{m-q-1}$ ).

The assumption that  $N$  has boundary is not essential in the Theorem and the Corollary. But these results are trivial for closed  $N$ : if  $N$  is closed and  $M \neq N$ , then the assumptions are never fulfilled because  $H^n(N) \neq H^n(M)$ . Note that the conclusion of the Theorem for closed manifolds is not always fulfilled, for there are closed manifolds non-embeddable in the same dimension as the corresponding punctured manifolds.

Now let us present motivation for the second corollary of the Theorem. From the construction of the map  $\tilde{f}$  above it follows that

$$\text{if } N \text{ embeds into } \mathbb{R}^m, \text{ then there exists an equivariant map } \tilde{N} \rightarrow S^{m-1}. \quad (*)$$

The existence of an equivariant map  $\tilde{N} \rightarrow S^{m-1}$  can be checked for many cases [4, beginning of §2], [6], [7, 1.7.1], [20], [1, 7.1]. Thus if a converse to  $(*)$  is true, the embedding problem is reduced to manageable (although not trivial) algebraic problems. So in 1960s there appeared a problem to find conditions under which the converse to  $(*)$  is true. The converse for  $(*)$  was known to be

*true for an  $n$ -polyhedron  $N$  and  $2m \geq 3n + 3$  or  $m = 2n = 2$*

[7,25,23], see also [17,18], [12, §4], [19].

*false for each pair  $(m, n)$  such that  $\max\{4, n\} \leq m \leq \frac{3n}{2} + 1$  and*

*some  $n$ -polyhedron  $N$  [10,9,15,5,21].*

In the only remaining cases  $m = 3$  and  $n \in \{2, 3\}$  it was unknown if the converse to  $(*)$  is true. The counterexamples to the converse of  $(*)$  for  $m = n \geq 4$  and  $m = n + 1 \geq 4$  [10,9] cannot be directly extended to  $m = 3$  because they used  $m$ -dimensional (Mazur) contractible manifolds distinct from the  $m$ -ball, which apparently do not exist for  $m = 3$ .

Recall that a *homology  $n$ -ball* is an  $n$ -manifold with boundary whose homology groups are the same as those of the  $n$ -ball.

**Proposition.** *The converse to (\*) is false in the cases  $m = 3$  and  $n \in \{2, 3\}$ : if  $N$  is either a non-trivial homology ball or its special spine, then  $N$  does not embed into  $\mathbb{R}^3$  but there exists an equivariant map  $\tilde{N} \rightarrow S^2$ .*

**Proof.** The non-embeddability follows because if a special spine of a homology ball  $N$  embeds into  $\mathbb{R}^3$ , then the regular neighborhood in  $\mathbb{R}^3$  of this spine is homeomorphic to  $N$  [3], which contradicts to the non-triviality of  $N$ .

It suffices to prove the existence of an equivariant map  $\tilde{N} \rightarrow S^2$  for a homology 3-ball  $N$ . For this case this existence follows from the Theorem because an inclusion of the standard ball into  $N$  induces an isomorphism in cohomology.

We also present an alternative proof of the existence of an equivariant map  $\tilde{N} \rightarrow S^2$  for a homology 3-ball  $N$ , which proof is shorter than that of Theorem. Analogously to [1, end of §7.1] (or by Steps 2 and 3 below) it suffices to prove that  $H^i(\tilde{N}; \mathbb{Z}) = 0$  for each  $i \geq 3$ . We prove this for  $i = 3$  (the proof for each  $i \geq 4$  is analogous). We omit  $\mathbb{Z}$ -coefficients from the notation. Let  $V$  be a closed regular neighborhood in  $N \times N$  of the diagonal. Then

$$\begin{aligned} H^3(\tilde{N}) &\cong H^3(N \times N - \mathring{V}) \cong H_3(N \times N - \mathring{V}, \partial(N \times N - \mathring{V})) \\ &\cong H_3(N \times N, V \cup \partial(N \times N)) \cong H_2(V \cup \partial(N \times N)) = 0. \end{aligned}$$

Here the first isomorphism follows because  $N \times N - \mathring{V}$  is a deformation retract of  $\tilde{N}$ , the second one by Lefschetz duality (recall that  $\tilde{N}$  is orientable if  $N$  a homology ball), the third one by excision, and the fourth one by exact sequence of pair. Using the Mayer–Vietoris sequence for

$$\partial(N \times N) = N \times \partial N \bigcup_{\partial N \times \partial N} \partial N \times N$$

and noting that  $\partial N \cong S^2$ , we prove that  $H_2(\partial(N \times N)) = 0$ . Using the Mayer–Vietoris sequence for  $V \cup \partial(N \times N)$  and noting that  $V \simeq N$  and  $V \cap \partial(N \times N)$  is a regular neighborhood in  $N \times N$  of the diagonal of  $\partial N$ , i.e. is homotopy equivalent to  $\partial N \cong S^2$ , we prove the last isomorphism.  $\square$

Another proof of the Proposition can be obtained by using the fact that for the homology 3-ball  $N$ , which is a punctured boundary of the Mazur 4-manifold, there exists an equivariant map  $\Sigma \tilde{N} \rightarrow S^3$  [11]. Indeed, one can show that the obstruction to equivariant desuspension of this map on  $\tilde{P}$  (where  $P$  is the special spine of  $N$ ) lies in  $H^4(\tilde{P}, \mathbb{Z})$ , which group is trivial because  $P$  is acyclic [24]. We do not present the details because we already have a complete proof of the Proposition.

**Proof of the Theorem** (without the ‘more precisely’ part). Denote  $N^* = \tilde{N}/(x, y) \sim (y, x)$ . Consider the bundle

$$g: \frac{\tilde{N} \times S^{m-1}}{(x, y, s) \sim (y, x, -s)} \xrightarrow{S^{m-1}} N^*, \quad \text{given by } g[(x, y, s)] = [(x, y)].$$

Equivariant maps  $\tilde{N} \rightarrow S^{m-1}$  up to equivariant homotopy are in 1–1 correspondence with cross-sections of  $g$  up to equivalence [4, beginning of §2], [6], [7, 1.7.1], [20], [1, 7.1]. So we need to prove that any section on  $M^*$  can be extended to a section on  $N^*$ , and such an extension is unique up to equivalence. This follows because below (in Steps 1, 2 and 3) we prove that

$$H^{i+1}(N^*, M^*; \pi_i(S^{m-1})_{tw}) \cong 0 \quad \text{and} \\ H^i(N^*, M^*; \pi_i(S^{m-1})_{tw}) \cong 0 \quad \text{for each } i.$$

Here the coefficients are twisted according to the double cover  $\tilde{N} \rightarrow N^*$  and the involution on  $\pi_i(S^{m-1})$  induced by the antipodal involution on  $S^{m-1}$  (so the twisting is trivial for  $m$  even).

*Step 1. Proof that  $H^i(\tilde{N}, \tilde{M}; G) \cong 0$  for each  $i$  and any non-twisted  $G$ .* Let  $N_0$  and  $M_0$  the interiors of  $N$  and  $M$ , respectively. The collaring theorem for the boundary of a manifold says that there is a neighborhood of  $\partial M$  in  $M$  which is homeomorphic to the product  $\partial M \times [0, 1]$  so that  $\partial M \times \{0\}$  is mapped homeomorphically to the boundary. Therefore there is an embedding of  $\phi: N \rightarrow N_0$  which is a homotopy inverse of the inclusion  $i: N_0 \rightarrow N$ . In a similar fashion the map  $\phi \times \phi: \tilde{N} \rightarrow \tilde{N}_0$  is the homotopy inverse of the inclusion  $\tilde{N}_0 \rightarrow \tilde{N}$ . Analogous observations hold for  $N$  replaced by  $M$ . So it suffices to prove Step 1 for  $N$  and  $M$  replaced by  $N_0$  and  $M_0$ .

Let  $x_0 \in M_0 \subset N_0$  be a base point for  $M_0$  and  $N_0$ . Consider the following mapping of bundles (which are given by projections onto the first factor):

$$\begin{array}{ccccc} M_0 - x_0 & \longrightarrow & \tilde{M}_0 & \longrightarrow & M_0 \\ \downarrow \subset & & \downarrow \subset & & \downarrow \subset \\ N_0 - x_0 & \longrightarrow & \tilde{N}_0 & \longrightarrow & N_0 \end{array}$$

Consider the spectral sequences associated to the bundles above. Observe that  $M_0 - x_0 \simeq M_0 \vee S^{n-1}$ . Hence the action of  $\pi_1(M_0)$  in the (co-)homology  $H^i(M_0 - x_0)$  is trivial for all  $i$  (the same holds for the second bundle, where  $M$  is replaced by  $N$ ). This follows for  $i = n$  because  $H^n(M_0 - x_0) = 0$  and for  $i < n - 1$  because the projection onto the first factor is the restriction of the trivial bundle  $M_0 \times M_0 \rightarrow M_0$ , while  $H^i(M_0 - x_0) \cong H^i(M_0)$ . For  $i = n - 1$  we have  $H^{n-1}(M_0 - x_0) \cong H^{n-1}(M_0) \oplus \mathbb{Z}$ . The action of an element  $\alpha \in \pi_1(M_0)$  is given by the identity on the first summand and multiplication by the sign of the loop on  $\mathbb{Z}$ . Since the manifolds are orientable, the action is identical.

Consider the restriction-induced homomorphism

$$j: H^p(N_0; H^q(N_0 - x_0; G)) \rightarrow H^p(M_0; H^q(M_0 - x_0; G))$$

of the  $E_2$ -terms of the Leray–Serre cohomology spectral sequences of the above bundles. By excision the inclusion of the pairs  $(M_0, M_0 - x_0) \rightarrow (N_0, N_0 - x_0)$  induces an isomorphism in cohomology. Applying 5-lemma for the inclusion-induced mapping of long exact sequences of these pairs we obtain that the inclusion  $M_0 - x_0 \rightarrow N_0 - x_0$  induces an isomorphism in cohomology. Hence using the triviality of the action and the Universal Coefficients Theorem we obtain that the restriction-induced homomorphism  $j$  is an isomorphism for all  $p$  and  $q$ . By the Zeeman Comparison Theorem of spectral sequences [26], the restriction  $H^i(\tilde{N}_0; G) \rightarrow H^i(\tilde{M}_0; G)$  is an isomorphism. This implies Step 1.

A statement on cohomology of compact manifolds should have a proof involving only cohomology of compact manifolds (recall that we may assume that  $\tilde{N} = \tilde{N}_\varepsilon$  is compact). The above proof has such an interpretation in terms of only compact spaces. Step 1 can also be proved analogously to proof of the Proposition above. We do not present details in order to spare space.

*Step 2. Proof that  $H^i(N^*, M^*; G) \cong 0$  for each  $i$  and any non-twisted coefficient system  $G$ .* Consider the double coverings  $\tilde{M} \rightarrow M^*$  and  $\tilde{N} \rightarrow N^*$ . The restriction induces a map of their Cartan–Leray spectral sequences, see [2, Chapter XVI, Theorem 8.4 on p. 354]. For the  $E_2$ -terms we have a map

$$E_2^{p,q}(N^*) = H^p(\mathbb{Z}_2, H^q(\tilde{N})) \rightarrow E_2^{p,q}(M^*) = H^p(\mathbb{Z}_2, H^q(\tilde{M})),$$

which is an isomorphism by Step 1 and the exact sequence of pair  $(\tilde{N}, \tilde{M})$ . Hence the restriction induces an isomorphism  $H^i(N^*; G) \rightarrow H^i(M^*; G)$ , and the result follows.

*Another proof of Step 2.* The  $\mathbb{Z}_2$  principal fibration  $\tilde{M} \rightarrow M^*$  is induced by the inclusion  $i: M^* \rightarrow N^*$  from the  $\mathbb{Z}_2$  principal fibration  $\tilde{N} \rightarrow N^*$ . The classifying map of the principal fibration  $\tilde{N} \rightarrow N^*$  is a map  $f: N^* \rightarrow \mathbb{R}P^\infty$ , and the classifying map for  $\tilde{M} \rightarrow M^*$  is the composition  $f \circ i$ . Therefore we obtain a commutative diagram of bundles

$$\begin{array}{ccccc} F_1 & \longrightarrow & E_1 & \longrightarrow & \mathbb{R}P^\infty \\ \downarrow \subset & & \downarrow \subset & & \downarrow = \\ F_2 & \longrightarrow & E_2 & \longrightarrow & \mathbb{R}P^\infty \end{array}$$

where  $F_1, E_1, F_2$  and  $E_2$  have the same homotopy type as  $\tilde{M}, M^*, \tilde{N}$  and  $N^*$ , respectively. Now Step 2 follows because for the  $E_2$ -term of cohomology spectral sequence of this pair of bundles we have  $E_2^{p,q} = H^p(\mathbb{R}P^\infty; H^q(E_2, E_1; G)) \cong 0$  for each  $p, q$  and any non-twisted  $G$ .

*Step 3. Proof that  $H^i(N^*, M^*; G_{tw}) \cong 0$  for each  $i$  and a local coefficient system  $G_{tw}$  associated to the double cover  $\tilde{N} \rightarrow N^*$  and certain involution  $\varphi: G \rightarrow G$  of a module  $G$ , which is a finitely-generated Abelian group.* It suffices to consider the case when  $G = \mathbb{Z}$ . The local system induced by  $G_{tw}$  on  $\pi_1(\tilde{N})$  is non-twisted. By naturality,  $G_{tw}$  induces a local system in  $M^*$ . Clearly, the local system over  $M^*$  also induces a trivial system on  $\tilde{M}$ , and the local system on  $M^*$  is trivial if and only if  $G_{tw}$  is trivial on  $N^*$ . Observe that there is only one non-trivial local system  $\mathbb{Z}_{tw}$  on  $N^*$  which induces the trivial local system on  $\tilde{N}$ , and the same for  $M^*$ . Now consider the map between the Smith–Richardson–Thom–Gysin sequences associated with the double covers  $\tilde{M} \rightarrow M^*$  and  $\tilde{N} \rightarrow N^*$  (see the Smith–Richardson–Thom–Gysin Sequence Theorem below, cf. [1, §7]).

$$\begin{array}{ccccccccc} H^j(N^*, \mathbb{Z}) & \longrightarrow & H^j(\tilde{N}, \mathbb{Z}) & \longrightarrow & H^j(N^*, \mathbb{Z}_{tw}) & \longrightarrow & H^{j+1}(N^*, \mathbb{Z}) & \longrightarrow & H^{j+1}(\tilde{N}, \mathbb{Z}) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ H^j(M^*, \mathbb{Z}) & \longrightarrow & H^j(\tilde{M}, \mathbb{Z}) & \longrightarrow & H^j(M^*, \mathbb{Z}_{tw}) & \longrightarrow & H^{j+1}(M^*, \mathbb{Z}) & \longrightarrow & H^{j+1}(\tilde{M}, \mathbb{Z}) \end{array}$$

Suppose by induction that  $H^{j+1}(N^*, \mathbb{Z}_{tw}) \rightarrow H^{j+1}(M^*, \mathbb{Z}_{tw})$  is an isomorphism. Certainly this is true for  $j+1$  big enough so we can start the induction process. Then the inductive step follows by Steps 1, 2 and the five lemma.

*The proof of the ‘more precisely’ part of the Theorem.* Now assume that the inclusion  $M \rightarrow N$  induces isomorphism in cohomology only down to dimension  $l$ . It suffices to prove, Steps 1, 2 and 3 for  $i \geq n + l$  instead of arbitrary  $i$ . We show how to do this for Step 1, and the required changes in the proofs of Steps 2 and 3 are obvious. The condition  $H^i(\tilde{N}, \tilde{M}; G) \cong 0$  for  $i \geq n + l$  is equivalent to  $H^i(\tilde{N}_0, \tilde{M}_0; G) \cong 0$  for  $i \geq n + l$ . Consider the spectral sequences associated to the bundles given in Step 1. Recall that

$$H^n(N_0 - x_0; G) = H^n(N_0; G) = H^n(M_0 - x_0; G) = H^n(M_0; G) = 0$$

for any coefficients  $G$ . As in Step 1 the action of  $\pi_1(M_0)$  in the (co-)homology  $H^i(M_0 - x_0)$  is trivial for all  $i$ . Analogous statement holds for the second fibration. By excision the inclusion of the pairs  $(M_0, M_0 - x_0) \rightarrow (N_0, N_0 - x_0)$  induces an isomorphism in cohomology. Applying the 5-lemma for the inclusion-induced mapping of long exact sequences of these pairs we obtain that the inclusion  $M_0 - x_0 \rightarrow N_0 - x_0$  induces an isomorphism in cohomology for  $i \geq l$ . Using all the facts above and the Universal Coefficient Theorem we see that the restriction induces an isomorphism

$$j: H^p(N_0; H^q(N_0 - x_0; G)) \rightarrow H^p(M_0; H^q(M_0 - x_0; G))$$

of the  $E_2$ -terms of the Leray–Serre cohomology spectral sequences of the above bundles for  $p + q \geq n + l - 1$ . So the restriction induces an isomorphism of  $E_2^{p,q}$  for  $p + q \geq n + l$  and an epimorphism for  $p + q = n + l - 1$ . Now using standard argument of homological algebra as in the Zeeman Comparison Theorem of spectral sequences [26] we obtain that the homomorphism between the terms  $E_r^{p,q}$  is an isomorphism for  $p + q \geq n + l$  and an epimorphism for  $p + q = n + l - 1$ . Since  $E_{n-l} = E_{n-l+1} = \dots = E_\infty$ , it follows that the restriction induces on  $E_\infty$  terms an isomorphism for  $p + q \geq n + l$  and an epimorphism for  $p + q = n + l - 1$ . Hence the restriction  $H^i(\tilde{N}; G) \rightarrow H^i(\tilde{M}; G)$  is an isomorphism for  $i \geq n + l$  and an epimorphism for  $p + q = n + l - 1$ . Therefore by the long exact sequence of pair it follows that  $H^i(\tilde{N}, \tilde{M}; G) \cong 0$  for  $i \geq n + l$ .  $\square$

For the reader’s convenience we present below the following slight extension of known Smith–Richardson–Thom–Gysin sequence.

**Smith–Richardson–Thom–Gysin Sequence Theorem.** *Let  $X$  be a connected space,  $X' \rightarrow X$  a double covering and  $G$  a module with an involution  $\phi$ . Consider the local coefficient system  $G_\phi$  on  $X$  associated to the double covering and the involution  $\phi$ , the system  $G_{-\phi}$  on  $X$  associated to double covering  $p$  and involution  $-\phi$ , and the trivial local system  $G$  on  $X'$ . Then there is a long exact sequence*

$$\begin{aligned} \dots \rightarrow H^{p-1}(X, G_\phi) \rightarrow H^p(X, G_{-\phi}) \rightarrow H^p(X', G) \\ \rightarrow H^p(X, G_\phi) \rightarrow H^{p+1}(X, G_{-\phi}) \rightarrow \dots \end{aligned}$$

*If 2 is invertible in  $G$  (in particular, if either  $G = \mathbb{Q}$  or  $G = \mathbb{Z}_p$  for  $p$  an odd prime), then we have splittable short exact sequence*

$$\begin{aligned} 0 \rightarrow H^p(X, G_{-\phi}) \rightarrow H^p(X', G) \rightarrow H^p(X, G_\phi) \rightarrow 0 \quad \text{so that} \\ H^p(X', G) \cong H^p(X, G_{-\phi}) \oplus H^p(X, G_\phi). \end{aligned}$$

If  $G = \mathbb{Z}$  and  $\phi = \text{id}$ , then we get long exact sequence

$$\begin{aligned} \cdots \rightarrow H^{p-1}(X, \mathbb{Z}) \rightarrow H^p(X, \mathbb{Z}_{tw}) \rightarrow H^p(X', \mathbb{Z}) \\ \rightarrow H^p(X, \mathbb{Z}) \rightarrow H^{p+1}(X, \mathbb{Z}_{tw}) \rightarrow \cdots \end{aligned}$$

If  $G = \mathbb{Z}$  and  $\phi = -\text{id}$ , then we get long exact sequence

$$\begin{aligned} \cdots \rightarrow H^{p-1}(X, \mathbb{Z}_{tw}) \rightarrow H^p(X, \mathbb{Z}) \rightarrow H^p(X', \mathbb{Z}) \\ \rightarrow H^p(X, \mathbb{Z}_{tw}) \rightarrow H^{p+1}(X, \mathbb{Z}) \rightarrow \cdots \end{aligned}$$

**Proof.** The given local systems  $\pi_1(X) \rightarrow \text{Aut}(G)$  on  $X$  factor through  $\mathbb{Z}_2$ , so for simplicity we will regard  $G$  as a  $\mathbb{Z}[\mathbb{Z}_2]$  module. Consider the fibration  $F \rightarrow X' \rightarrow X$  which is the double covering, where  $F$  is a two-points set. For the spectral sequence with local coefficients [16] we have  $E_2^{p,q} = H^p(X, H^q(F, G))$ . But these groups are trivial for each  $q > 0$ . Then the spectral sequence contains at most one non-vanishing line. Hence

$$H^p(X', G_\phi) \cong E_\infty^{p,0} \cong E_2^{p,0} \cong H^p(X, (G \oplus G)_{tw}),$$

where the local coefficient system  $(G \oplus G)_{tw}$  is associated to the double covering and the involution  $\phi \oplus \phi$ . Namely, the local coefficient system is defined by the automorphism  $\tau$  of  $H^0(F, G) \cong G \oplus G$  induced by the map  $F \rightarrow F$  permuting the coordinates and by the homomorphism  $\phi \oplus \phi$ . This homomorphism  $\tau$  is given by  $\tau(a, b) = (\phi(b), \phi(a))$ .

Now let  $H = \{(m, -m) \in G \oplus G \mid m \in G\}$ . The subgroup  $H$  is invariant under the action of  $\mathbb{Z}[\mathbb{Z}_2]$  on  $G \oplus G$ , so  $H$  is a sub-module. The involution on this submodule is given by  $\tau(m, -m) = (\phi(-m), \phi(m)) = (-\phi(m), \phi(m))$ . Hence  $H$  as a module is isomorphic to  $G$  with the involution  $-\phi$ , i.e. to  $G_{-\phi}$ . The quotient  $(G \oplus G)/G_\phi$  is also a  $\mathbb{Z}[\mathbb{Z}_2]$  module and it is easy to see that it is isomorphic to  $G_\phi$ . Now the first part of the Smith–Richardson–Thom–Gysin Sequence Theorem follows from the long exact sequence in cohomology associated with the short exact sequence of coefficients of  $\mathbb{Z}[\mathbb{Z}_2]$ -modules.

The further part where 2 is invertible follows from the fact that the short exact sequence of  $\mathbb{Z}[\mathbb{Z}_2]$  modules

$$0 \rightarrow H \rightarrow G \oplus G \rightarrow (G \oplus G)/H \rightarrow 0$$

splits. Namely, the  $\mathbb{Z}[\mathbb{Z}_2]$  homomorphism  $s : (G \oplus G)/H \cong G \rightarrow G \oplus G$  defined by  $s(m) = 1/2(m, m)$  is a splitting. The part where  $G = \mathbb{Z}$  is clear.  $\square$

## Acknowledgements

This work was initiated during the visit of the first author to Faculty of Mechanics and Mathematics of Lomonosov Moscow State University in May 2002, and the first author thanks Department of Differential Geometry and Applications of this Faculty for hospitality during his stay. These results were first presented by the second author at the Seminar on low-dimensional topology and group theory (University of Frankfurt), Topological seminar (University of Ljubljana) and Workshop on geometric topology and dimension theory (Mathematical Institute of Polish AS) in November and December 2002, and the second author thanks these institutions for their invitations. We would like to acknowledge A.Yu. Volovikov for useful discussions and the referee for his remarks.



## References

- [1] M. Adachi, *Embeddings and Immersions*, Transl. Math. Monographs, vol. 124, Amer. Math. Soc., Providence, RI, 1993.
- [2] H. Cartan, S. Eilenberg, *Homological Algebra*, Oxford University Press, Princeton University Press, Oxford–Princeton, 1956.
- [3] B.G. Casler, An embedding theorem for connected 3-manifolds with boundary, *Proc. Amer. Math. Soc.* 16 (1965) 556–559.
- [4] P.E. Conner, E.E. Floyd, Fixed points free involutions and equivariant maps, *Bull. Amer. Math. Soc.* 66 (1960) 416–441.
- [5] M.H. Freedman, V.S. Krushkal, P. Teichner, Van Kampen’s embedding obstruction is incomplete for 2-complexes in  $\mathbb{R}^4$ , *Math. Res. Lett.* 1 (1994) 167–176.
- [6] A. Haefliger, Plongements de variétés dans le domaine stable, *Seminare Bourbaki* 245 (1962).
- [7] A. Haefliger, Plongements différentiables dans le domaine stable, *Comment. Math. Helv.* 37 (1962–63) 155–176.
- [8] K. Horvatič, Embedding manifolds with low-dimensional spine, *Glas. Mat.* 4(24) (1) (1969) 101–116.
- [9] L.S. Husch,  $\varepsilon$ -maps and embeddings, in: *General Topological Relations to Modern Analysis and Algebra*, VI, Heldermann, Berlin, 1988, pp. 273–280.
- [10] S. Mardešić, J. Segal,  $\varepsilon$ -mappings and generalized manifolds, *Michigan Math. J.* 14 (1967) 171–182.
- [11] J. Malešić, D. Repovš, A. Skopenkov, Embeddings into  $\mathbb{R}^m$  and the deleted product obstruction, *Bol. Soc. Mat. Mexicana* 9 (2003) 165–170.
- [12] D. Repovš, A. Skopenkov, New results on embeddings of polyhedra and manifolds into Euclidean spaces, *Uspekhi Mat. Nauk* 54 (6) (1999) 61–109, in Russian; English transl.: *Russian Math. Surveys* 54 (6) (1999) 1149–1196.
- [13] C.P. Rourke, B.J. Sanderson, *Introduction to Piecewise-Linear Topology*, *Ergeb. Math.*, vol. 69, Springer-Verlag, Berlin, 1972.
- [14] M. Scharlemann, Isotopy and cobordism of homology spheres in spheres, *J. London Math. Soc.*, Ser. 2 16 (3) (1977) 559–567.
- [15] J. Segal, S. Spiez, Quasi-embeddings and embedding of polyhedra in  $\mathbb{R}^m$ , *Topology Appl.* 45 (1992) 275–282.
- [16] J. Siegel, Higher order cohomology operations in local coefficient theory, *Amer. J. Math.* 89 (1967) 909–931.
- [17] A.B. Skopenkov, On the deleted product criterion for embeddability of manifolds in  $\mathbb{R}^m$ , *Comment. Math. Helv.* 72 (1997) 543–555.
- [18] A.B. Skopenkov, On the deleted product criterion for embeddability in  $\mathbb{R}^m$ , *Proc. Amer. Math. Soc.* 126:8 (1998) 2467–2476.
- [19] A.B. Skopenkov, On the Haefliger–Hirsch–Wu invariants for embeddings and immersions, *Comment. Math. Helv.* 77 (2002) 78–124.
- [20] A. Skopenkov, *Embedding and Knotting of Higher-Dimensional Manifolds in Euclidean Spaces*, *London Math. Soc. Lect. Notes*, submitted for publication.
- [21] J. Segal, A. Skopenkov, S. Spiez, Embedding of polyhedra in  $\mathbb{R}^m$  and the deleted product obstruction, *Topology Appl.* 85 (1998) 335–344.
- [22] C.T.C. Wall, *Theory of Handle Decompositions*, *Differential Topology*, vol. IV, Cambridge University Press, Cambridge, 1964 (mimeographed notes).
- [23] C. Weber, Plongements de polyèdres dans le domaine métastable, *Comment. Math. Helv.* 42 (1967) 1–27.
- [24] C. Weber, Deux remarques sur les plongements d’un AR dans un espace euclidien, *Bull. Acad. Polon. Sci. Ser. Sci. Math. Astronom. Phys.* 16 (1968) 851–855.
- [25] W.T. Wu, *A Theory of Embeddings, Isotopy and Immersions of Polyhedra in Euclidean Space*, Science Press, Peking, 1965.
- [26] E.C. Zeeman, A proof of the comparison theorem for spectral sequence, *Proc. Cambridge Philos. Soc.* 53 (1957) 57–62.